

Brownian motion approach to the ideal gas of relativistic particles

Ryszard Zygadlo

Marian Smoluchowski Institute of Physics, Jagiellonian University, Reymonta 4, PL-30059 Kraków, Poland

(Dated: February 2, 2008)

The relativistic generalization of a free Brownian motion theory is presented. The global characteristics of the relaxation are *explicitly* found for the velocity and momentum (stochastic) kinetics. It is shown that the thermal corrections, to the both relaxation times T (of stationary autocorrelations) and transient relaxation time of momentum, appear slowing down the processes. The transient relaxation time of the velocity does not depend *explicitly* on temperature, $T(v_0) = m(v_0)/\gamma \equiv \varepsilon_0/\gamma c^2$, and it is proportional to the initial energy of a relativistic Brownian particle.

During the last twenty years the Brownian motion approach has become the most important tool for discovering and studying the spectacular phenomena, the *critical slowing down* [1], *noise-induced transitions* [2], *stochastic resonance* [3], *ratchet dynamics* [4], and *resonant activation* [5], in (nonlinear) nonequilibrium systems. In the most of thermodynamical applications the nonlinear Langevin equation appears as an overdamped limit of a dissipative dynamic in a certain external potential [6]. The important conclusion of the present paper is that, within the Markovian diffusion theory, the nonlinear stochastic kinetic results already on the basic level of (spatially homogeneous) ideal gases description if the relativistic particles are considered. Particularly the relativistic free Brownian motion theory based on identical general assumption as the original one [i.e., including the interaction with the thermal bath by a *systematic force*, the dissipative kinematic friction proportional to the velocity $-\gamma v$, and an additive *random force*, the thermal fluctuation, given by Gaussian white noise ξ_t]

$$\dot{p}_t = -\gamma v + \xi_t, \quad (1)$$

where [$c = 1$ is used throughout the paper] $p = mv/\sqrt{1-v^2}$ is a relativistic momentum of a Brownian particle, $v = p/\sqrt{p^2 + m^2} = p/\varepsilon$, $\varepsilon = \sqrt{p^2 + m^2}$, and $\langle \xi_t \xi_s \rangle = 2D\delta(t-s)$, is consistent both with the thermodynamical requirements of the equilibrium Gibbs ensemble theory and with the special relativity requirements of transformational properties of the momentum probability density distribution.

First note that the general Kramers–Fokker–Planck kinetic equation (see, e.g., the equation (26.26) of the Ref. [7]) of a Markovian diffusion description simplifies to the form

$$\partial_t P = \partial_p L_{22}[(\partial_p H_1)P + k_B T \partial_p P] \quad (2)$$

for the spatially homogeneous system of free noninteracting particles [of the Hamiltonian $H_1(p)$]. The kinetic coefficient $L_{22} = L_{22}(p, T)$ is in principle determined by the properties of the interaction with the thermal bath and do not depend on p for the potential interaction. For the latter case (of the state independent diffusion) the Langevin description Eq. (1), and the corresponding Fokker–Planck one [8], Eq. (2), are equivalent in view of the identification $L_{22} = \gamma = \gamma(T)$,

$D = \gamma k_B T$ (the Einstein relation, or the *fluctuation-dissipation theorem*), and $\partial_p H_1 = v$. The Eq. (2) is the most general form of the diffusion in the momentum space leading to the Gibbs–Boltzmann stationary state $W(p) \propto e^{-\beta H_1(p)}$ of thermal equilibrium ($\beta = 1/k_B T$). The corresponding (nonlinearly coupled) Langevin equation (in the Stratonovich interpretation [8]) is

$$\dot{p}_t = -\gamma g^2 v + D g g' + g \xi_t, \quad (3)$$

where $L_{22} = \gamma g^2$ and $g' = \partial g / \partial p$.

The (any) corresponding probability density distributions $W(p)$ and $W'(p')$ in a resting and moving (with a constant velocity V) reference frame are related by non-covariant formula $\varepsilon W(p) = \varepsilon' W'(p')$, where ε and p should be expressed as functions of ε' and p' , according to the Lorentz transformation [9]. Because the time increments (counted from the initial preparation of the system) are given by t' and $t = t'/\sqrt{1-V^2}$, respectively, the relation

$$P'(p', t') = \frac{1 + V v'}{\sqrt{1 - V^2}} P\left(\frac{p' + V \varepsilon'}{\sqrt{1 - V^2}}, \frac{t'}{\sqrt{1 - V^2}}\right) \quad (4)$$

is required. One has $v = (v' + V)/(1 + V v')$, $\xi_t = (1 - V^2)^{1/4} \xi_{t'}$, and, using the Stratonovich calculus, $dp/dt = (1 + V v') dp'/dt'$, thus the transformed stochastic kinetic (1) takes the form

$$\frac{dp'}{dt'} = -\gamma \frac{v' + V}{(1 + V v')^2} + \frac{(1 - V^2)^{1/4}}{1 + V v'} \xi_{t'}, \quad (5)$$

where $v' = p'/\sqrt{p'^2 + m^2}$. It is easy to verify that the r.h.s. of (4) solves the Fokker–Planck equation corresponding to the Eq. (5) if (and only if) $P(p, t)$ satisfies the proper Eq. (2) related to the kinetic (1). The same conclusion applies also to the general case (3).

The Eq. (1), $\dot{p}_t = -\gamma p/\sqrt{p^2 + m^2} + \xi_t$, for the momentum or, equivalently, the (Stratonovich) equation for the velocity,

$$\dot{v}_t = -\gamma v(1 - v^2)^{3/2}/m + \xi_t(1 - v^2)^{3/2}/m, \quad (6)$$

thus provides the relativistic generalization of the diffusion Markovian description of equilibration for noninteracting free particles system [10]. The normalized stationary distributions are

$$W(p) = \frac{1}{2mK_1(m\gamma/D)} e^{-(\gamma/D)\sqrt{p^2 + m^2}} \quad (7)$$

$$\text{and} \quad W(v) = \frac{(1-v^2)^{-3/2}}{2K_1(m\gamma/D)} e^{-(\gamma/D)m/\sqrt{1-v^2}}, \quad (8)$$

where $\gamma = D\beta$. K_n is a modified Bessel function. The second moments read

$$\langle p^2 \rangle = \frac{m}{\beta} \frac{K_2(m\beta)}{K_1(m\beta)}, \quad \langle v^2 \rangle = 1 - \int_{m\beta}^{\infty} dz \frac{K_0(z)}{K_1(m\beta)}, \quad (9)$$

so the (even) moments of the velocity, as incomplete integrals of Bessel functions, are not given in closed analytical form.

The nonlinear kinetic (1) or (6) does not belong to the class of the known solvable models and the time-dependent solution is not known. Nevertheless, there are still the quantities, useful to characterize the relaxation of the nonlinear systems, which can be expressed by *quadratures* of the (*necessarily autonomous*) drift and diffusion coefficients, without solving the kinetic equation. These are the complete (over time) integrals of (various) transient (single-time) moments [11] and stationary (two-times) correlation functions [12]. In fact, making use of $P(x, t|x_0) = e^{t\hat{L}}\delta(x-x_0)$, where $\hat{L}(x_0)$ is a *backward* Fokker-Planck operator [8], and computing $X(x_0) = \int_0^\infty dt \langle x_t \rangle_{x_0}$, where the integrand is a conditional moment equal x_0 at the initial time $t=0$, by a formal integration $\int_0^\infty dt e^{t\hat{L}} = -\hat{L}^{-1}$ one obtains

$$\hat{L}X = -x_0, \quad (10)$$

which is a nonhomogeneous linear equation of the first order for the function dX/dx_0 . The $\langle x \rangle_{st} = 0$ is required for the convergence of X and the condition $X(0) = 0$ determines a first constant of integration for Eq. (10). The second one should be chosen in such way that the correct (deterministic) result is recovered when the noise strength $D \rightarrow 0$. The Eq. (10) differs from the Pontryagin equation for the mean first passage time [13] in nonhomogeneous term ($-x_0$ instead of -1). The integrals of stationary autocorrelations are then given by $S_x = \int_0^\infty dt \langle x_t x_0 \rangle_{st} = \int dx W(x) X(x)x$. The last quantity (i.e., the power spectrum function taken at zero frequency $S_x = S_x(0)$), is called, after some normalization (usually divided by the stationary variance) resulting in the proper (time) dimension, the *relaxation time* T [12]. The analytical expression for this quantity has been first obtained by Jung and Risken [14]. Then, by a similar consideration, the formulae for the quantities of the former type, called generally nonlinear relaxation times, have been found [15, 11].

Identifying the appropriate coefficients from Eqs. (1) and (6), and carrying out the integrals of the Jung and Risken formula [14] we obtain the exact results

$$S_v = D/\gamma^2, \quad (11)$$

$$S_p = Dm^2/\gamma^2 + \frac{3mD^2}{\gamma^3} \frac{K_2(m\gamma/D)}{K_1(m\gamma/D)} - D^3/\gamma^4. \quad (12)$$

The same Eq. (11) is obtained for the ordinary Brownian motion, $S_p^o = m^2 S_v^o = Dm^2/\gamma^2$. Introducing $T_x = S_x/\langle x^2 \rangle$ and *using m and m/γ as units of momentum and time* [which is equivalent with considering the generic forms of Eqs. (1), (6) with a single (*implicit*) parameter $D = \tau$ and $m = \gamma = 1$] one gets for *dimensionless* quantities

$$T_v = \left\{ \int_0^1 dv \frac{\exp[(-\tau\sqrt{1-v^2})^{-1}]}{K_1(\tau^{-1})} \right\}^{-1}, \quad (13)$$

$$T_p = \frac{1 - \tau^2 + 3\tau K_2(\tau^{-1})/K_1(\tau^{-1})}{K_2(\tau^{-1})/K_1(\tau^{-1})}, \quad (14)$$

where τ is a *dimensionless temperature*, $\tau = D/m\gamma = k_B T/mc^2$. Note that (13) is the alternative form of $T_v = \tau/\langle v^2 \rangle$, where $\langle v^2 \rangle$ is given by Eq. (9), which is more suitable for numerical computations. The respective quantities of the ordinary Brownian motion are simply equal to the inverse of the relaxation rate of a linear model (1), $T_v^o = T_p^o = 1$, and do not depend *explicitly* on temperature [16]. In contrast both T_v or T_p increases with temperature, see Fig. 1, approaching the asymptotic $T_v(\tau) \approx \tau + \pi/2$ or $T_p(\tau) \approx (5/2)\tau$, respectively. The former asymptotic law follows from representation (9) [the complete integral of K_0 is equal to $\pi/2$], the latter is obtained by the use of the asymptotic formula $K_n(1/\tau) \approx (n-1)!(2\tau)^n/2$ (for $\tau \rightarrow \infty$) [17] in Eq. (14). For small $\tau \rightarrow 0$ $T_v(\tau) \approx 1 + (3/2)\tau$ and $T_p(\tau) \approx 1 + 3\tau$. So, the decay of both (*stationary*) autocorrelations in a relativistic case proceed slower in higher temperatures, and slower than for the ordinary Brownian motion. We want to stress however that, in the case of nonlinear system, the relaxation time T_x should *not* be interpreted as a characteristic of the equilibration process from a certain *nonstationary* state. One has in particular $T_x(\tau=0) = 1$, which would be misunderstood that the (deterministic) evolution of the relativistic and classical system proceeds with the same (or at least similar) speed. Meanwhile it may be generally proved that at the limit $\tau \rightarrow 0$ only the lowest order terms of the drift and diffusion coefficients contribute to T_x . Thus, such a conclusion applies at most to the relaxation from the close to the equilibrium states.

In order to compare the nonstationary behavior let us define the *transient relaxation time*

$$T_\tau(x_0) = x_0^{-1} \int_0^\infty dt \langle x_t \rangle_{x_0}, \quad (15)$$

which is related to the quantity X , Eq. (10), via $x_0 T_\tau(x_0) = X_\tau(x_0)$. Then $P_\tau(p_0)$ satisfies

$$-p_0 P'_\tau / \sqrt{1+p_0^2} + \tau P''_\tau = -p_0.$$

The particular solution of the nonhomogeneous equation is $P'_\tau = \sqrt{1+p_0^2} + \tau$, whereas the (general) solution of

the homogeneous equation is singular at $\tau = 0$. Thus,

$$T_\tau(p_0) = [p_0\sqrt{1+p_0^2} + \log(p_0 + \sqrt{1+p_0^2})]/2p_0 + \tau. \quad (16)$$

Similarly,

$$[-v_0(1-v_0^2)^{3/2} - 3\tau v_0(1-v_0^2)^2]V'_\tau + \tau(1-v_0^2)^3V''_\tau = -v_0$$

has a (regular at $\tau = 0$) particular solution $V'_\tau = (1-v_0^2)^{-3/2}$, so

$$T_\tau(v_0) = 1/\sqrt{1-v_0^2} \equiv \varepsilon_0. \quad (17)$$

The deterministic (generic) Stokes equations, $\dot{p} = -p/\sqrt{p^2+1}$ and $\dot{v} = -v(1-v^2)^{3/2}$, are solvable by quadratures. The solution, with the initial condition x_0 , is

$$t(x) = t_x(x) - t_x(x_0), \quad (18)$$

where $t_p(p) = -\sqrt{1+p^2} + \log|(1+\sqrt{1+p^2})/p|$, and $t_v(v) = -(1-v^2)^{-1/2} + \log|(1+\sqrt{1-v^2})/v|$, respectively. One can verify that the direct calculation of $\int_0^\infty dt x(t)/x_0 = \int_{x_0}^0 dx t(x)/x_0$ yields $T_0(x_0)$ as given by Eq. (16) or (17). The previous results (11) and (12) are also recovered when $T_\tau(x_0)$ is integrated with $x_0^2 W(x_0)$, where W is a stationary probability density distribution (8) or (7), respectively. Both (dimensionless) transient relaxation times for nonrelativistic Brownian motion are equal to unity. The relativistic relaxation of momentum depends both on the temperature and the initial state. $T_\tau(p_0)$, Eq. (16), is an increasing function of both arguments and has the following asymptotic $T_\tau(p_0) \approx 1 + \tau + p_0^2/6$, for $p_0 \rightarrow 0$, and $T_\tau(p_0) \approx |p_0|/2 + \tau$, for $p_0 \rightarrow \infty$. The result for the velocity is somehow unexpected. The Eq. (17) shows that indeed the relativistic relaxation proceeds slower, however does not depend on temperature. It turns out that with the increase of temperature the early stage of evolution becomes faster in such a way that compensates the subsequent slower long-time behavior, see Fig. 3. The transient relaxation time $T_\tau(v_0)$ is equal to the (dimensionless) initial energy of the relativistic Brownian particle ε_0 . The plots of $T_0(x_0)$ are shown in Fig. 2.

The equation (1) (in a generic form) has been solved numerically by the Runge–Kutta method for 10^4 sample realizations of the white Gaussian noise, for different initial conditions and temperatures. $\langle p_t \rangle$ and $\langle v_t \rangle$ have then been computed as the arithmetical average of appropriate values obtained for different trajectories of the noise. The typical curves are plotted in Figs. 3 and 4, together with the corresponding deterministic results (18). The solution of the relativistic Stokes equation for velocity changes qualitatively, when v_0 exceeds $1/2$. In fact below this value of the initial state the function $v(t)$ is convex down for all t [as well as (always) $p(t)$]. However for $v_0 > 1/2$ at the early stage of evolution the $v(t)$ appears convex up [in order to achieve the required asymptote

$v(t) = 1 = c$ for $v_0 \rightarrow 1$]. The former *subrelativistic* case, with particular $v_0 = 2/5$ ($p_0 \approx 0.436$), and the latter, with $p_0 = 4$ ($v_0 \approx 0.970$), are presented.

The results of the paper may be summarized as follows. We have shown that using the relations of the relativistic dynamics we can generalize the Brownian motion theory in a way consistent with thermodynamical requirements. The general solvable equation (10) for the quantities simply related to the transient relaxation times (15) has been found. These, as well the stationary relaxation times T , have been found analytically for the relativistic momentum and velocity (stochastic) kinetics, Eqs. (13, 14, 16, 17). All these quantities (considered as dimensionless) are equal unity for the ordinary Brownian motion. The relaxation times $T_x(\tau)$, by the normalization equal unity in the deterministic limit $\tau \rightarrow 0$, exhibit positive first order corrections for small temperatures, becoming just proportional to τ in a high temperature asymptotic. Thus the decay of correlations in a stationary state proceed slower than for the nonrelativistic Brownian motion. The transient relaxation time [describing the relaxation from a given initial state x_0 , and by the normalization equal unity for the deterministic system set initially at the equilibrium, $T_0(0) = 1$] of momentum $T_\tau(p_0)$ is an affinic function of temperature. The asymptotic behavior for the large argument(s) reads $T_\tau(p_0) \approx |p_0|/2 + \tau$. The transient relaxation time for the velocity (17) does not depend on temperature, and it is equal to the dimensionless initial energy of the relativistic Brownian particle. The *thermal corrections* to the considered global characteristics of the relaxation are linear with the slope of the unity order and appear (even) for the noninteracting free relativistic Brownian particles system. The corrections can be observed for sufficiently high *physical* temperatures. For, e.g., hydrogen particle (proton) the dimensionless quantity, which characterize the deviation from the ordinary nonrelativistic behavior, $\tau = k_B T/mc^2 \approx 10^{-13} T[\text{K}]$. So for the temperature $T \sim 1\text{MeV} \sim 10^{10}\text{K}$ the *relative* correction is of the order 0.1%. The correction to transient relaxation time of the velocity has the kinematic origin only and for $T_\tau(p_0)$ both kinds of corrections are decoupled. Note that the general dependence on temperature in physical units, introduced (multiplicatively) by the coefficient $\gamma(T)$, should be counted separately [16].

The three-dimensional generalization of the proposed diffusion description is obvious, however in contrast to the ordinary Brownian motion theory it leads to the set of the three coupled ($\vec{v} = \vec{p}/\varepsilon = \vec{p}/\sqrt{m^2 + \vec{p}^2}$) Langevin equations (with statistically independent noises in each direction) or equivalently to the (forward) Fokker–Planck description [8] $[\nabla = \partial/\partial \vec{p}]$

$$\partial_t P(\vec{p}, t|\vec{p}_0) = \gamma \nabla \circ (\vec{p} P/\varepsilon) + D \nabla^2 P. \quad (19)$$

The required form of the (Gibbs–Boltzmann) stationary distribution is still recovered, however the calculations of relaxation times cannot be analytically completed in this case.

-
- [1] A. Schenzle and H. Brand, Phys. Rev. A **20**, 1628 (1979); R. Graham and A. Schenzle, *ibid.* **25**, 1731 (1982); M. Suzuki, K. Kaneko, and S. Takesue, Prog. Theor. Phys. **67**, 1676 (1982); M. Suzuki, S. Takesue, and F. Sasagawa, *ibid.* **68**, 98 (1982).
- [2] W. Horsthemke and R. Lefever, *Noise-Induced Transitions*, (Springer, Berlin, 1984).
- [3] L. Gammaitoni, P. Haenggi, P. Jung, and F. Marchesoni, Rev. Mod. Phys. **70**, 223 (1998).
- [4] P. Reimann, Phys. Rep. **361**, 57 (2002).
- [5] C.R. Doering and J.C. Gadoua, Phys. Rev. Lett. **69**, 2318 (1992); P. Pechukas and P. Haenggi, *ibid.* **73**, 2772 (1994); P. Reimann, *ibid.* **74**, 4576 (1995); P. Reimann and T.C. Elston *ibid.* **77**, 5328 (1996).
- [6] see, e.g., *Noise in Nonlinear Dynamical Systems* Vol. 1, Eds. F. Moss and P.V.E. McClintock (University, Cambridge, 1989).
- [7] D.N. Zubarev, *Nonequilibrium Statistical Thermodynamics*, (Plenum, New York, 1974).
- [8] C.W. Gardiner, *Handbook of Stochastic Methods* (Springer, Berlin, 1983); H. Risken, *The Fokker-Planck Equation: Methods of Solution and Application* (Springer, Berlin, 1983).
- [9] L. Landau and E. Lifshitz, *The Classical Theory of Fields* (Pergamon, Oxford, 1971).
- [10] The different general approach to the subject, based on the Boltzmann kinetic equation, is presented by S.R. de Groot, W.A. van Leeuwen, and Ch.G. van Weert, *Relativistic Kinetic Theory* (North-Holland, Amsterdam, 1980).
- [11] P. Colet, M. San Miguel, J. Casademunt, and J.M. Sancho, Phys. Rev. A **39**, 149 (1989).
- [12] J.M. Sancho, M. San Miguel, S.L. Katz, and J.D. Gunton, Phys. Rev. A **26**, 1589 (1982); S. Faetti, C. Festa, L. Fronzoni, P. Grigolini, F. Marchesoni, and V. Palleschi, Phys. Lett. A **99**, 25 (1983).
- [13] L. Pontryagin, A. Andronov, and A. Vitt, Zh. Eksp. Teor. Fiz. **3**, 165 (1933); translated in *Noise in Nonlinear Dynamical Systems* Vol. 1, Eds. F. Moss and P.V.E. McClintock (University, Cambridge, 1989).
- [14] P. Jung and H. Risken, Z. Phys. B **59**, 469 (1985); J.M. Sancho, R. Mannella, P.V.E. McClintock, and F. Moss Phys. Rev. A **32**, 3639 (1985).
- [15] J.I. Jimenez-Aquino, J. Casademunt, and J.M. Sancho, Phys. Lett. A **133**, 364 (1988); J. Casademunt, J.I. Jimenez-Aquino, and J.M. Sancho, Physica A **156**, 628 (1989).
- [16] All the remaining consideration concern on dimensionless quantities when particularly the time unit is fixed by m/γ . Thus the changes of values of quantities in physical units introduced by $\gamma = \gamma(T)$ are not included in our formulations claiming the temperature dependence. The temperature dependence of $\gamma = \gamma(T)$ is determined by the properties of the interaction with the thermal bath. For the hard sphere model one has in particular $\gamma(T) \propto \sqrt{T}$ (which corresponds to the ordinary nonrelativistic result), for $\tau \equiv k_B T/mc^2 \ll 1$ and $\gamma(T) \propto T$, for high temperatures $\tau \gg 1$, see Ref. [10].
- [17] I.M. Gradshteyn and I.M. Ryzhik, *Table of Integrals, Series and Products* (Academic, New York, 1965)

FIG. 1: Plots of T_p [Eq. (14), upper curve] and T_v [Eq. (13)] vs τ (dimensionless temperature). The latter has been computed numerically by the Romberg method.

FIG. 2: Plots of $T_0(p_0)$ [Eq. (16), $\tau = 0$, upper curve] vs p_0 and $T_\tau(v_0)$ [Eq. (17)] vs $10 \times v_0$.

FIG. 3: Plots of $\langle v_t \rangle_{v_0}/v_0$ vs t . The upper and lower curves correspond to $v_0 \approx 0.970$ and $v_0 = 2/5$; respectively. The deterministic solutions, Eq. (18), are marked by 0 ($\tau = 0$). The dotted curve, e^{-t} , corresponds to the nonrelativistic universal result. The remaining curves have been obtained numerically for $\tau = 0.2$ and 0.6 , respectively.

FIG. 4: The same for $\langle p_t \rangle_{p_0}/p_0$ vs t .







